


A REFINEMENT OF SARKOVSKII'S THEOREM(U) ARMY BALLISTIC 1/1  
RESEARCH LAB ABERDEEN PROVING GROUND MD  
N P BHATIA ET AL. SEP 86 BRL-TR-2762


F/G 19/4 NL

[illegible]



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

12



US ARMY  
MATERIEL  
COMMAND

AD

TECHNICAL REPORT BRL-TR-2762

AD-A174 030

# A REFINEMENT OF SARKOVSKI'S THEOREM

Nam P. Bhatia  
Walter O. Egerland

September 1986

DTIC  
ELECTE  
NOV 14 1986  
S E D

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

US ARMY BALLISTIC RESEARCH LABORATORY  
ABERDEEN PROVING GROUND, MARYLAND

DTIC FILE COPY

86 11 14 068

Destroy this report when it is no longer needed.  
Do not return it to the originator.

Additional copies of this report may be obtained  
from the National Technical Information Service,  
U. S. Department of Commerce, Springfield, Virginia  
22161.

The findings in this report are not to be construed as an official  
Department of the Army position, unless so designated by other  
authorized documents.

The use of trade names or manufacturers' names in this report  
does not constitute indorsement of any commercial product.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

# TABLE OF CONTENTS

I.	INTRODUCTION	Page 5
II.	ELEMENTARY LEMMAS	Page 6
III.	PRINCIPAL RESULTS	Page 7
IV.	REMARKS	Page 11
	REFERENCES	Page 13
	DISTRIBUTION LIST	Page 15

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



# I. INTRODUCTION

Let  $f: R \rightarrow R$  be continuous and  $x_0 \in R$ . The orbit of  $x_0$  under  $f$  is defined as the set  $\{x: x = f^n(x_0), n = 0, 1, \dots\}$ , where, for every positive integer  $n$ ,  $f^n$  is the  $n$ -th iterate of  $f$ ,  $f^1 = f$ , and  $f^0(x_0) = x_0$ . We shall write  $x_n := f^n(x_0)$  for a given  $x_0 \in R$  and call  $x_1, x_2, \dots$  the successors of  $x_0$ . A pre-orbit of a given  $x_0 \in R$  is any (finite or infinite) sequence  $x_0, x_{-1}, x_{-2}, \dots$  such that  $f(x_{-n}) = x_{-(n-1)}$  for all  $n$  for which  $x_{-n}$  is defined. The points  $x_{-1}, x_{-2}, \dots$  in any such sequence are called predecessors of  $x_0$ . A point  $c_0$  is called critical if  $f(c_0) = c_0$ , i.e., a critical point of  $f$  is a fixed point of  $f$ . A periodic point  $x_0$  of period  $p > 1$  ( $p$  a positive integer) is a point for which the relations  $f^p(x_0) = x_0$ ,  $f^k(x_0) \neq x_0$ ,  $1 \leq k < p$ , hold. If  $x_0$  is a periodic point of period  $p$ , its orbit is denoted by  $(x_0, x_1, \dots, x_{p-1})$ . We shall denote the  $k$ th iterate of  $x_0$  under the function  $f^m$  by  $x_k^m$ ,  $k = 0, 1, \dots$ . Thus  $x_k^m := (f^m)^k(x_0) = x_{mk}$ , and, in particular,  $x_0^m = x_k^0 = x_0$  for all nonnegative integers  $k$  and  $m$ .

Definition. Let  $f: R \rightarrow R$  be continuous and  $x_0 \in R$ .  $f$  has a loop of order  $n$  if  $x_0$  has a pre-orbit  $(x_0, x_{-1}, \dots, x_{-n})$  such that either

$$x_0 < x_{-n} < x_{-(n-1)} < \dots < x_{-2} < x_{-1}$$

or

$$x_0 > x_{-n} > x_{-(n-1)} > \dots > x_{-2} > x_{-1}.$$

$f$  has an infinite loop if  $x_0$  has an infinite pre-orbit  $(x_0, x_{-1}, \dots, x_{-n}, \dots)$  such that either

$$x_0 < \dots < x_{-n} < x_{-(n-1)} < \dots < x_{-2} < x_{-1}$$

or

$$x_0 > \dots > x_{-n} > x_{-(n-1)} > \dots > x_{-2} > x_{-1}.$$

A loop of order  $(n - 1)$  is called an  $n$ -periodic loop if  $x_0 = x_{-n}$ .

We adopt the following concise notation: we say property  $P(k)$  holds if  $f$  has a periodic orbit of period  $k$ . Thus  $P(1)$ ,  $L(k)$ ,  $L(=)$  mean that  $f$  has a critical point, a periodic loop of period  $k$ , an infinite loop, respectively. Similarly,  $P^n(k)$ ,  $L^n(k)$ ,  $L^n(=)$  shall mean that  $f^n$  has a  $k$ -periodic orbit,  $k$ -periodic loop, an infinite loop, respectively.

In this notation, Sarkovskii's theorem and our refinement read as follows.

**Theorem (Sarkovskii).** Let  $f: R \rightarrow R$  be continuous. Then

$$\begin{aligned} P(3) &\Rightarrow P(5) \Rightarrow P(7) \Rightarrow \dots \Rightarrow \\ P(2 \cdot 3) &\Rightarrow P(2 \cdot 5) \Rightarrow P(2 \cdot 7) \Rightarrow \dots \Rightarrow \\ P(2^2 \cdot 3) &\Rightarrow P(2^2 \cdot 5) \Rightarrow P(2^2 \cdot 7) \Rightarrow \dots \Rightarrow \\ \dots &\Rightarrow \\ P(2^3) &\Rightarrow P(2^2) \Rightarrow P(2) \Rightarrow P(1). \end{aligned}$$

**Theorem (SR).** Let  $f: R \rightarrow R$  be continuous. Then

$$\begin{aligned} L(\infty) &\dots \Rightarrow L(5) \Rightarrow L(4) \Rightarrow L(3) \Leftarrow \\ P(3) &\Rightarrow P(5) \Rightarrow P(7) \Rightarrow \dots \Rightarrow \\ L^2(\infty) &\dots \Rightarrow L^2(5) \Rightarrow L^2(4) \Rightarrow L^2(3) \Leftarrow \\ P(2 \cdot 3) &\Rightarrow P(2 \cdot 5) \Rightarrow P(2 \cdot 7) \Rightarrow \dots \\ L^{2^2}(\infty) &\dots \Rightarrow L^{2^2}(5) \Rightarrow L^{2^2}(4) \Rightarrow L^{2^2}(3) \Leftarrow \\ P(2^2 \cdot 3) &\Rightarrow P(2^2 \cdot 5) \Rightarrow P(2^2 \cdot 7) \Rightarrow \dots \\ \dots &\dots \\ \dots &\Rightarrow P(2^3) \Rightarrow P(2^2) \Rightarrow P(2) \Rightarrow P(1). \end{aligned}$$

A. N. Sarkovskii obtained the fundamental result which bears now his name in his seminal paper<sup>1</sup> of 1964. The purpose of this paper is to prove Theorem (SR).

## II. ELEMENTARY LEMMAS

It follows from the definition of a periodic loop that every three-periodic orbit is a three-periodic loop and that an  $(n + 1)$ -periodic loop implies the existence of a loop of order  $n$ .

**Lemma 2.1.** If  $f$  has a critical point  $c_0$  such that  $c_0 < c_{-2} < c_{-1}$ ,  $f$  has an infinite loop satisfying

$$c_0 < \dots < c_{-n} < \dots < c_{-2} < c_{-1}.$$

The same statement holds with all inequalities reversed.

**Proof.** Since  $f(c_{-2}) = c_{-1}$  and  $c_0 < c_{-1}$ , there exists  $c_{-3} \in (c_0, c_{-2})$ . Repeating this argument establishes the lemma.

**Lemma 2.2.** If  $f$  has a critical point  $c_0$  such that  $c_{-1} < c_{-3} < c_0 < c_{-2}$ ,  $c_0$  has

<sup>1</sup> A. N. Sarkovskii, "Coexistence of cycles of a continuous map of a line into itself," Ukrain. Mat. Zh. 16 (1964), 61-71.



an infinite pre-orbit satisfying

$$c_{-1} < c_{-3} < \dots < c_0 < \dots < c_{-4} < c_{-2}.$$

In particular,  $f^2$  has two infinite loops. The same statement holds with all inequalities reversed.

Proof. Since  $f(c_{-2}) = c_{-1} < c_{-3}$  and  $c_0 > c_{-3}$ , there exists  $c_{-4} \in (c_0, c_{-2})$ , and since  $f(c_{-3}) = c_{-2} > c_{-4} > c_0$ , there exists  $c_{-5} \in (c_{-3}, c_0)$ . Repeating this argument proves the lemma.

Lemma 2.3.  $P^{2^k}(n) \iff P(2^k \cdot n)$ ,  $n, k = 1, 2, \dots$ .

Proof. It suffices to show that  $P^2(n) \iff P(2 \cdot n)$ . If  $(x_0, x_1, \dots, x_{2n-1})$  is a  $2n$ -periodic orbit of  $f$ ,  $(x_0^2, x_1^2, \dots, x_{n-1}^2)$  is an  $n$ -periodic orbit of  $f^2$ .

Hence  $P(2 \cdot n) \implies P^2(n)$ . If  $(x_0^2, x_1^2, \dots, x_{n-1}^2)$  is an  $n$ -periodic orbit of  $f^2$ , we consider the set  $\{x_0, x_1, \dots, x_{2n-1}\}$ , where  $x_{2n} = x_n^2 = x_0^2 = x_0$ . If  $x_0 \neq x_k$ ,  $k=1, 2, \dots, 2n-1$ , then  $C = (x_0, x_1, \dots, x_{2n-1})$  is a  $2n$ -periodic orbit of  $f$ . Otherwise, there is a smallest odd  $k$ ,  $1 < k < 2n$ , such that  $x_0 = x_k$ , i.e.,  $x_0$  is an odd-periodic point of  $f$ . But then, by Sarkovskii's theorem,  $f$  has periodic orbits of every even period and, therefore, in particular, a  $2n$ -periodic orbit. Hence  $P^2(n) \implies P(2 \cdot n)$  and the proof of the lemma is complete.

### III. PRINCIPAL RESULTS

Let  $C = (x_0, x_1, \dots, x_{n-1})$  be any  $n$ -periodic orbit of  $f$ . We define the subsets

$$\begin{aligned} C^+ &= \{x_i \in C: x_{i+1} > x_i\}, \quad C^- = \{x_i \in C: x_{i+1} < x_i\}, \\ D^+ &= \{x_i \in C: x_{i+2} > x_{i+1} > x_i\}, \quad D^- = \{x_i \in C: x_{i+2} < x_{i+1} < x_i\}. \end{aligned}$$

The sets  $C^+$  and  $C^-$  are non-empty since  $\min C \in C^+$  and  $\max C \in C^-$ . Letting further  $a_0^+ = \min C^+ (= \min C)$ ,  $b_0^+ = \max C^+$ ,  $a_0^- = \min C^-$ , and  $b_0^- = \max C^- (= \max C)$ , it is clear that either  $a_0^+ \leq b_0^+ < a_0^- \leq b_0^-$  or  $a_0^+ < a_0^- < b_0^+ < b_0^-$ .

Theorem 3.1. If  $a_0^+ \leq b_0^+ < a_0^- \leq b_0^-$  and  $D^+ \cup D^- \neq \emptyset$ ,  $f$  has a critical point  $c_0$  such that  $f^2$  has two infinite loops  $(d_0^2, d_{-1}^2, d_{-2}^2, \dots)$  and  $(c_0^2, c_{-1}^2, c_{-2}^2, \dots)$  satisfying

$$d_{-1}^2 < d_{-2}^2 < \dots < d_0^2 = c_0 = c_0^2 < \dots < c_{-2}^2 < c_{-1}^2.$$

In particular,  $L^2(*)$  holds.

Proof. It is sufficient to assume that  $D^+ \neq \emptyset$ . Then, if we let  $\beta_0^+ = \max D^+$ , we have

$$a_0^+ \leq \beta_0^+ < \beta_1^+ \leq b_0^+ < a_0^- \leq \beta_2^+ \leq b_0^-.$$

and conclude the existence of a critical point  $c_0$  and a predecessor  $c_{-1}$  such that

$$a_0^+ \leq \beta_0^+ < c_{-1} < \beta_1^+ \leq b_0^+ < c_0 < a_0^- \leq b_0^-.$$

We consider now the set  $E^- = \{x_i \in C^- : x_{i+1} < c_{-1}\}$ .  $E^-$  is non-empty since  $a_{n-1}^+ \in C^-$  and  $a_n^+ = a_0^+ < c_{-1}$ . Letting  $r_0^- = \min E^-$ , we have

$$a_0^+ \leq r_1^- < c_{-1} < b_0^+ < c_0 < a_0^- \leq r_0^- \leq b_0^-.$$

This shows that, since  $c_0 > c_{-1}$  and  $r_1^- < c_{-1}$ , there exists a predecessor  $c_{-2}$  such that

$$a_0^+ < c_{-1} < b_0^+ < c_0 < c_{-2} < r_0^- \leq b_0^-.$$

Our construction implies that

$$(i) \text{ if } x_i \in C^+ \text{ and } x_i > c_{-1}, \text{ then } x_{i+1} \in C^-$$

$$(ii) \text{ if } x_i \in C^- \text{ and } x_i < r_0^-, \text{ then } x_{i+1} > c_{-1}.$$

Hence, there is an  $x_i \in C^+$ ,  $x_i > c_{-1}$  such that  $x_{i+1} \geq r_0^-$ . For otherwise we would have  $b_i^+ \in (c_{-1}, r_0^-)$  for all  $i$ , contradicting the fact that  $C$  is the orbit of  $b_0^+$  (thus  $b_i^+ = a_0^+$  for some  $i > 1$ ). We now choose  $\delta_0^+ \in C^+$  such that  $\delta_0^+ > c_{-1}$  and  $\delta_1^+ \geq r_0^-$  to obtain

$$a_0^+ < c_{-1} < \delta_0^+ \leq b_0^+ < c_0 < c_{-2} < r_0^- \leq \delta_1^+ \leq b_0^-.$$

But this implies that we may choose a predecessor  $c_{-3}$  in the interval  $(c_{-1}, \delta_0^+)$ , and hence that  $c_0$  and its predecessors  $c_{-1}$ ,  $c_{-2}$ , and  $c_{-3}$  satisfy the

inequality  $c_{-1} < c_{-3} < c_0 < c_{-2}$ . Appeal to Lemma 2.2 completes the proof.

**Theorem 3.2.** If  $a_0^+ < a_0^- < b_0^+ < b_0^-$ , there exist critical points  $d_0, c_0$  of  $f$  and two infinite loops  $(d_0, d_{-1}, d_{-2}, \dots)$  and  $(c_0, c_{-1}, c_{-2}, \dots)$  of  $f$  satisfying

$$d_{-1} < d_{-2} < \dots < d_0 \leq c_0 < \dots < c_{-2} < c_{-1}.$$

In particular,  $L(*)$  holds.

**Proof.** We note first that there is  $\alpha_0^- \in C^-$  and  $\beta_0^+ \in C^+$  such that

- (i)  $a_0^+ < a_0^- \leq \alpha_0^- < \beta_0^+ \leq b_0^+ < b_0^-$
- (ii) if  $x_i \in C$ , then  $x_i \leq \alpha_0^-$  or  $x_i \geq \beta_0^+$
- (iii) if  $x_i \in C$  and  $\alpha_0^- < x_i \leq b_0^+$ , then  $x_i \in C^+$
- (iv) if  $x_i \in C$  and  $b_0^+ < x_i \leq b_0^-$ , then  $x_i \in C^-$ .

We now show that there are predecessors  $c_{-1}$  and  $c_{-2}$  of the critical point  $c_0 \in (\alpha_0^-, \beta_0^+)$  that satisfy the inequality  $c_0 < c_{-2} < c_{-1}$ . The set  $A^- = \{x_i \in C^- : x_i > b_0^+ \text{ and } x_{i+1} \leq \alpha_0^-\}$  is non-empty (otherwise  $\beta_n^+ \geq \beta_0^+$  for all integers  $n \geq 0$ , a contradiction). Let  $r_0^- = \min A^-$ . We have  $r_0^- > b_0^+$  and observe that the set  $A^+ = \{x_i \in C^+ : \beta_0^+ \leq x_i \leq b_0^+ \text{ and } x_{i+1} \geq r_0^-\}$  is non-empty (since otherwise  $\beta_n^+$  will satisfy  $\beta_0^+ \leq \beta_n^+ < r_0^-$  for  $n \geq 0$ , a contradiction). We choose any  $y_0^+ \in A^+$  and have

$$\alpha_0^- < c_0 < \beta_0^+ \leq y_0^+ \leq b_0^+ < r_0^- \leq b_0^-.$$

Hence

$$c_0 < c_{-2} < y_0^+ \leq b_0^+ < c_{-1} < r_0^- \leq b_0^-,$$

where the existence of  $c_{-1}$  follows from  $b_1^+ > c_0$  and  $r_1^- < c_0$  and that of  $c_{-2}$  from  $c_0 < c_{-1}$  and  $y_1^+ \geq r_0^- > c_{-1}$ . The infinite loop  $(c_0, c_{-1}, c_{-2}, \dots)$  satisfying  $c_0 < \dots < c_{-2} < c_{-1}$  follows from Lemma 2.1. An analogous procedure locates a critical point  $d_0$  and predecessors  $d_{-1}, d_{-2}$  such that  $d_{-1} < d_{-2} < d_0 \leq c_0$ , and hence an infinite loop  $(d_0, d_{-1}, d_{-2}, \dots)$  satisfying  $d_{-1} < d_{-2} < \dots < d_0 \leq$

$c_0$ . This completes the proof.

**Theorem 3.3.** If  $f$  has a loop of order  $n \geq 3$ ,  $f$  has two distinct  $n$ -periodic loops. In particular,  $L(n)$  holds.

**Proof.** Let  $(x_0, x_{-1}, \dots, x_{-n})$  be a loop of order  $n \geq 3$  of  $f$  such that

$$x_0 < x_{-n} < \dots < x_{-2} < x_{-1}.$$

Since there is a critical point  $c_0 \in (x_{-2}, x_{-1})$ , there are predecessors  $c_{-1}, c_{-2}, \dots, c_{-(n-2)}$  such that

$$x_0 < x_{-n} < c_{-(n-2)} < x_{-(n-1)} < \dots < c_{-1} < x_{-2} < c_0 < x_{-1}.$$

We consider now the set

$$S = \{y_0 \in \mathbb{R} : y_n < y_0 < c_{-(n-2)} < y_1 < \dots < y_{n-3} < c_{-1} < y_{n-2} < c_0 < y_{n-1}\}.$$

The set  $S$  is non-empty since  $x_{-n} \in S$  and open since  $f$  is continuous. Let  $(a_0, b_0)$  be the component of  $S$  such that  $x_{-n} \in (a_0, b_0)$ . Since  $c_{-(n-2)} \notin S$  and  $y_0 \in (a_0, b_0)$  implies  $y_0 < c_{-(n-2)}$ , we must have

$$- \infty < a_0 < b_0 < c_{-(n-2)}.$$

We first note that  $a_0 > -\infty$ . This is so because for every  $y_0 \in (a_0, b_0)$  we have  $y_n < y_0$  and  $y_n \in f^2([c_{-1}, c_0])$ , which is a compact set. Thus  $y_0 \geq \min f^2([c_{-1}, c_0]) > -\infty$ . This implies  $a_0 \geq \min f^2([c_{-1}, c_0]) > -\infty$ . Since  $a_0, b_0 \notin S$  and  $y_0 \in (a_0, b_0)$  implies  $y_0 \in S$ , we conclude by the continuity of  $f$  that

$$a_n = a_0 < c_{-(n-2)} < a_1 < \dots < a_{n-3} < c_{-1} < a_{n-2} < c_0 < a_{n-1}$$

and

$$b_n = b_0 < c_{-(n-2)} < b_1 < \dots < b_{n-3} < c_{-1} < b_{n-2} < c_0 < b_{n-1}.$$

Hence both  $a_0$  and  $b_0$  are  $n$ -periodic points, and since  $a_0 < b_0 < b_1 < \dots < b_{n-1}$ , the orbits of  $a_0$  and  $b_0$  are distinct. This completes the proof of the theorem.

**Corollary 3.3.** If  $f$  has an  $(n+1)$ -periodic loop,  $n \geq 3$ ,  $f$  has two distinct  $n$ -periodic loops.

**Theorem 3.4.**  $L(\infty) \implies \dots \implies L(4) \implies L(3) \implies P(5) \implies P(7) \implies \dots \implies L^2(\infty)$ .

Proof. If  $f$  has an infinite loop satisfying

$$x_0 < \dots < x_{-2} < x_{-1},$$

the subset  $\{x_0, x_{-1}, \dots, x_{-n}\}$ ,  $n \geq 3$ , satisfies

$$x_0 < x_{-n} < \dots < x_{-2} < x_{-1},$$

and is, therefore, a loop of order  $n$ . By Theorem 3.3,  $L(n)$  holds. Hence

$L(\infty) \Rightarrow L(n)$ . By Corollary 3.3,  $L(n) \Rightarrow L(n-1)$ . The implications

$L(3) \Rightarrow P(5) \Rightarrow P(7) \Rightarrow \dots$  follow from Sarkovskii's theorem. Finally, to prove the implication  $P(2n+1) \Rightarrow L^2(\infty)$  for every  $n \geq 1$ , we note that if  $C = (x_0, x_1, \dots, x_{2n})$  is a  $(2n+1)$ -periodic orbit, then  $n(C^+) \neq n(C^-)$ , so that the hypothesis of either Theorem 3.1 or Theorem 3.2 is satisfied. In the first case  $L^2(\infty)$  holds by Theorem 3.1. In the second case, Theorem 3.2 implies that  $L(\infty)$  holds, and hence that  $L(3)$  holds. Now for any three-periodic orbit, the hypothesis of Theorem 3.1 holds trivially. Hence  $L^2(\infty)$  holds. This completes the proof.

Corollary 3.4.  $L^{2^k}(\infty) \dots \Rightarrow L^{2^k}(5) \Rightarrow L^{2^k}(3) \Rightarrow P(2^k \cdot 5) \Rightarrow P(2^k \cdot 7) \Rightarrow \dots \Rightarrow L^{2^{k+1}}(\infty)$ .

Proof. This follows from Theorem 3.4 and Lemma 2.3.

Proof of Theorem (SR). Theorem (SR) follows by combining Theorem 3.4, Corollary 3.4, Lemma 2.3 and Sarkovskii's theorem.

#### IV. REMARKS

1. Theorem (SR) is a step in the direction of obtaining a complete refinement of Sarkovskii's theorem that takes into account the orbit types of each period  $n$ . A periodic loop is only one of the orbit types of a given period. That certain orbit types imply the existence of infinite loops is implicit in Theorem 3.2 and is strikingly illustrated by the example  $f(x) = ax(1 - |x|)$ .  $f$  has the four-periodic orbit  $\left[\frac{1}{2}, \frac{a}{4}, \frac{1}{2}, -\frac{a}{4}\right]$ , where  $a \approx 4.411138875$  is given by

$$a^{-1} = \frac{1}{2} \left[ \left( \frac{1}{2} + \frac{1}{2} \sqrt{\frac{59}{27}} \right)^{1/3} + \left( \frac{1}{2} - \frac{1}{2} \sqrt{\frac{59}{27}} \right)^{1/3} \right]$$

This orbit satisfies the conditions of Theorem 3.2 and hence guarantees the existence of two infinite loops as well as two periodic loops of each period  $n \geq 3$  by Theorem (SR). The results of<sup>5</sup> ensure for this example merely the existence of a three-periodic orbit.

2. The results in this paper offer a novel approach to detecting chaos. Most practical methods for detecting chaos rely, either implicitly or explicitly, on the existence of odd periodic orbits<sup>2,3,4,5</sup>. However, lemmas 2.1 and 2.2 can be used by finding only a few predecessors of a critical point. Lemma 2.2, in particular, is independent of odd periods. A notable illustration is the example  $f(x) = x^2 - s$ , for which at

$$s^* = \frac{1}{3} \left[ 2 + \left( 3\sqrt{33} + 17 \right)^{1/3} - \left( 3\sqrt{33} - 17 \right)^{1/3} \right]$$

$L^2(\bullet)$  holds, with no odd period  $\neq 1$  being present.

<sup>2</sup> Gyorgy Targonski, "Topics in Iteration Theory," Studia Mathematica, Skript 6, Vandenhock and Ruprecht, Göttingen and Zürich (1981).

<sup>3</sup> Nam P. Bhatia and Walter O. Egerland, "On the Existence of Li-Yorke Points in the Theory of Chaos," Mathematics Research Report No. 84-4, Department of Mathematics, UMBC, July 1984. (To appear in Vol. 9, No. 10 of Nonlinear Analysis.)

<sup>4</sup> Nam P. Bhatia and Walter O. Egerland, "Non-periodic conditions for Chaos and Snap-Back Repellers," Transactions of the Second Army Conference on Applied Mathematics and Computing, ARO Report 85-1 159-164 (Also Mathematics Research Report No. 84-5, Department of Mathematics, UMBC, September 1984).

<sup>5</sup> Tien-Yien Li, Michal Misiurewicz, Giulio Pianigiani and James A. Yorke, "No Division Implies Chaos," Transactions of the American Mathematical Society, 273(1) (1982), 191-199.

## REFERENCES

- [1] A. N. Sarkovskii, "Coexistence of cycles of a continuous map of a line into itself," Ukrain. Mat. Zh. 16 (1964), 61-71.
- [2] Gyorgy Targonski, "Topics in Iteration Theory," Studia Mathematica, Skript 6, Vandenhoeck and Ruprecht, Göttingen and Zürich (1981).
- [3] Nam P. Bhatia and Walter O. Egerland, "On the Existence of Li-Yorke Points in the Theory of Chaos," Mathematics Research Report No. 84-4, Department of Mathematics, UMBC, July 1984. (To appear in Vol. 9, No. 10 of Nonlinear Analysis.)
- [4] Nam P. Bhatia and Walter O. Egerland, "Non-periodic conditions for Chaos and Snap-Back Repellers," Transactions of the Second Army Conference on Applied Mathematics and Computing, ARO Report 85-1, 159-164 (Also Mathematics Research Report No. 84-5, Department of Mathematics, UMBC, September 1984).
- [5] Tien-Yien Li, Michal Misiurewicz, Giulio Pianigiani and James A. Yorke, "No Division Implies Chaos," Transactions of the American Mathematical Society, 273(1) (1982), 191-199.

<u>Copies</u>	<u>Organization</u>	<u>Copies</u>	<u>Organization</u>
12	Administrator Defense Technical Info Center ATTN: DTIC-DDA Cameron Station Alexandria, VA 22304-6145	1	Commander U.S. Army Communications- Electronics Command ATTN: AMSEL-ED Fort Monmouth, NJ 07703
1	HQDA DAMA-ART-M Washington, D.C. 20310	1	Commander ERADCOM Technical Library ATTN: DELSD-L (Reports Section) Fort Monmouth, NJ 07703-5301
1	Commander U.S. Army Materiel Command ATTN: AMCDRA-ST 5001 Eisenhower Avenue Alexandria, VA 22333-0001	1	Commander U.S. Army Missile Command Research, Development & Engin- eering Center ATTN: AMSMI-RD Redstone Arsenal, AL 35898
1	Commander Armament R&D Center U.S. Army AMCCOM ATTN: SMCAR-TSS Dover, NJ 07801	1	Director U.S. Army Missile & Space Intelligence Center ATTN: AIAMS-YDL Redstone Arsenal, AL 35898-5500
1	Commander Armament R&D Center U.S. Army AMCCOM ATTN: SMCAR-TDC Dover, NJ 07801	1	Commander U.S. Army Tank Automotive Cmd ATTN: AMSTA-TSL Warren, MI 48397-5000
1	Director Benet Weapons Laboratory Armament R&D Center U.S. Army AMCCOM ATTN: SMCAR-LCB-TL Watervliet, NY 12189	1	Director U.S. Army TRADOC Systems Analysis Activity ATTN: ATAA-SL White Sands Missile Range, NM 88002
1	Commander U.S. Army Armament, Munitions and Chemical Command ATTN: SMCAR-ESP-L Rock Island, IL 61299	1	Commandant U.S. Army Infantry School ATTN: ATSH-CD-CSO-OR Fort Benning, GA 31905
1	Commander U.S. Army Aviation Research and Development Command ATTN: AMSAV-E 4300 Goodfellow Blvd St. Louis, MO 63120	1	Commander U.S. Army Development and Employ- ment Agency ATTN: MODE-TED-SAB Fort Lewis, WA 98433
1	Director U.S. Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035	1	AFWL/SUL Kirtland AFB, NM 87117
		1	Air Force Armament Laboratory ATTN: AFATL/DLODL Eglin AFB, FL 32542-5000



CopiesOrganization

10 Central Intelligence Agency  
Office of Central Reference  
Dissemination Branch  
Room GE-47 HQS  
Washington, D.C. 20502

ABERDEEN PROVING GROUND

Dir, USAMSAA  
ATTN: AMXSY-D  
AMXSY-MP, H. Cohen

Cdr, USATECOM  
ATTN: AMSTE-TO-F

Cdr, CRDC, AMCCOM  
ATTN: SMCCR-RSP-A  
SMCCR-MU  
SMCCR-SPS-IL

# USER EVALUATION SHEET/CHANGE OF ADDRESS

This Laboratory undertakes a continuing effort to improve the quality of the reports it publishes. Your comments/answers to the items/questions below will aid us in our efforts.

1. BRL Report Number \_\_\_\_\_ Date of Report \_\_\_\_\_

2. Date Report Received \_\_\_\_\_

3. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

4. How specifically, is the report being used? (Information source, design data, procedure, source of ideas, etc.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

5. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided or efficiencies achieved, etc? If so, please elaborate. \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

6. General Comments. What do you think should be changed to improve future reports? (Indicate changes to organization, technical content, format, etc.) \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

CURRENT ADDRESS      Name \_\_\_\_\_  
                                 Organization \_\_\_\_\_  
                                 Address \_\_\_\_\_  
                                 City, State, Zip \_\_\_\_\_

7. If indicating a Change of Address or Address Correction, please provide the New or Correct Address in Block 6 above and the Old or Incorrect address below.

OLD ADDRESS      Name \_\_\_\_\_  
                                 Organization \_\_\_\_\_  
                                 Address \_\_\_\_\_  
                                 City, State, Zip \_\_\_\_\_

(Remove this sheet along the perforation, fold as indicated, staple or tape closed, and mail.)

----- FOLD HERE -----

Director  
U.S. Army Ballistic Research Laboratory  
ATTN: SLCBR-DD-T  
Aberdeen Proving Ground, MD 21005-5066




NO POSTAGE  
NECESSARY  
IF MAILED  
IN THE  
UNITED STATES

OFFICIAL BUSINESS  
PENALTY FOR PRIVATE USE, \$300

**BUSINESS REPLY MAIL**  
FIRST CLASS PERMIT NO 12062 WASHINGTON, DC  
POSTAGE WILL BE PAID BY DEPARTMENT OF THE ARMY

Director  
U.S. Army Ballistic Research Laboratory  
ATTN: SLCBR-DD-T  
Aberdeen Proving Ground, MD 21005-9989



----- FOLD HERE -----

END

12-86

DTIC